



# MiR-574-5P, miR-1827, and miR-4429 as Potential Biomarkers for Schizophrenia

Omran Davarinejad<sup>1</sup> · Sajad Najafi<sup>2</sup> · Hossein Zhaleh<sup>3</sup> · Farzaneh Golmohammadi<sup>1</sup> · Farnaz Radmehr<sup>1</sup> · Mostafa Alikhani<sup>1</sup> · Reza Heidari Moghadam<sup>4</sup> · Yazdan Rahmati<sup>1</sup>

Received: 27 August 2021 / Accepted: 6 November 2021

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

## Abstract

Schizophrenia is a severe chronic debilitating disorder with millions of affected individuals. Diagnosis is based on clinical presentations, which are made when the progressive disease has appeared. Early diagnosis may help improve the clinical outcomes and response to treatments. Lack of a reliable molecular diagnostic invokes the identification of novel biomarkers. To elucidate the molecular basis of the disease, in this study we used two mRNA expression arrays, including GSE93987 and GSE38485, and one miRNA array, GSE54914, and meta-analysis was conducted for evaluation of mRNA expression arrays via metaDE package. Using WGCNA package, we performed network analysis for both mRNA expression arrays separately. Then, we constructed protein–protein interaction network for significant modules. Limma package was employed to analyze the miRNA array for identification of dysregulated miRNAs (DEMs). Using genes of significant modules and DEMs, a mRNA–miRNA network was constructed and hub genes and miRNAs were identified. To confirm the dysregulated genes, expression values were evaluated through available datasets including GSE62333, GSE93987, and GSE38485. The ability of the detected hub miRNAs to discriminate schizophrenia from healthy controls was evaluated by assessing the receiver-operating curve. Finally, the expression levels of genes and miRNAs were evaluated in 40 schizophrenia patients compared with healthy controls via Real-Time PCR. The results confirmed dysregulation of hsa-miR-574-5P, hsa-miR-1827, hsa-miR-4429, CREBRF, ARPP19, TGFBR2, and YWHAZ in blood samples of schizophrenia patients. In conclusion, three miRNAs including hsa-miR-574-5P, hsa-miR-1827, and hsa-miR-4429 are suggested as potential biomarkers for diagnosis of schizophrenia.

**Keywords** Schizophrenia · metaDE · WGCNA · microRNA · Real-Time PCR

✉ Yazdan Rahmati  
yazdanrahmati68@gmail.com

Sajad Najafi  
sajadnajafi1990@yahoo.com

<sup>1</sup> Clinical Research Development Center, Imam Khomeini and Mohammad Kermanshahi and Farabi Hospitals, Kermanshah University of Medical Sciences, Kermanshah, Iran

<sup>2</sup> Student Research Committee, Department of Medical Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup> Substance Abuse Prevention Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

<sup>4</sup> Cardiovascular Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

## Introduction

Schizophrenia (SCZ) is a chronic debilitating mental disease with a complex identity considered as one of the most mysterious human disorders (van Os and Kapur 2009a). It mainly appears during adulthood primary years affecting both genders, any race, and genetic backgrounds. According to the global estimations, about 20 million people are living with SCZ around the world (James et al. 2018). The affected patients are expected to have a reduced lifespan of 12–15 years compared to healthy individuals which show higher mortality compared to physical diseases (van Os and Kapur 2009b). SCZ diagnosis is based on psychopathological features including psychosis, hallucination or delusion, and impaired emotions and volition in absence of any medical disorder (Taylor et al. 1974). Thus, disease diagnosis is made when the clinical manifestations have appeared,

and thus, any diagnostic approach to screen the individuals at risk suggests special importance. The impairment of neurotransmitters like dopamine and gamma aminobutyric acid (GABA) has been the center of the main therapeutical strategies in SCZ; however, the results of current treatments are not satisfying indicating requirement for new concepts. Although the etiology of SCZ is still unknown likely to be multifactorial including genetic and environmental risk factors (Richetto and Meyer 2021), we already have known that individuals who have a history of SCZ in their first-grade family members have a higher risk of developing the disorder indicating high degrees of heritability (Lichtenstein et al. 2009). Furthermore, accumulating evidence suggests familial predisposition as the main risk factor for the disease in addition to other provoking conditions like pregnancy and simultaneous infections, impairments in neurological development, and cannabis abuse (Mäki et al. 2005). Different epigenetic mechanisms including DNA methylation, histone modifications, and regulation by non-coding RNAs (ncRNAs) also have been found to be involved in the SCZ pathogeny considered as molecular scars of environmental exposures (Richetto and Meyer 2021; Kuehner et al. 2019). They are particularly known to play role in neural development, and transcriptome studies show methylation changes and dysregulation of several epigenetic factors such as histone deacetylases (HDACs) and microRNAs (miRNAs) in brain samples of SCZ patients (Richetto and Meyer 2021; Sharma et al. 2008; Du et al. 2019). Additionally, recently a brilliant study has showed that transplantation of exosomes from SCZ patients mimics SCZ behaviors in mice (Du et al. 2021). Recent developments have suggested further consideration of epigenetic regulation in pathogeny and diagnostic and therapeutical potentials for SCZ in the future (Richetto and Meyer 2021).

MiRNAs are a novel group of ncRNAs with an average length of 22 nucleotides not encoding any protein unlike the known class of protein-coding messenger RNAs (mRNAs). Their functions have not been yet completely elucidated; however, numerous key regulatory roles in the eukaryotic gene expression have been described for an increasing number of identified miRNAs. In a variety of biological processes such as stress responses (Leung and Sharp 2010), cell proliferation, differentiation, and death (Bushati and Cohen 2007), embryogenesis, organ development, and function (Kloosterman and Plasterk 2006; Maes et al. 2008), miRNAs have been identified with regulatory functions in multicellular organisms. They mainly act at the post-transcriptional level in interaction with protein complexes via binding to complementary regions mostly located on the 3' untranslated region (UTR) of the target sponged mRNA and thus, drive their degradation by RNA-induced silencing complex (RISC) leading to repressing the expression of critical proteins which play role in essential cellular processes. As

expected, biogenesis, processing, nucleolar export, and stability of miRNAs are faced with precise regulation (Treiber et al. 2012) which also suggests their substantial roles. Constantly growing evidence has demonstrated the association of aberrant expression of miRNAs and numerous human diseases such as diabetes (Tang et al. 2008), osteoporosis (Van Wijnen et al. 2013), neurodegenerative diseases (Nelson et al. 2008), viral infections (Islam et al. 2019), and various types of cancer (Seven et al. 2014). Several miRNAs also have been shown to be aberrantly expressed in serum samples (Shi et al. 2012), cortex tissues (Perkins et al. 2007), and gene analyses of SCZ patients (Hauberg et al. 2016). As a result, differentially expressed miRNAs potentially can be considered as SCZ biomarkers (He et al. 2017). This approach can be beneficial particularly since the miRNAs are easily detected in blood samples of the patients and eventually can be employed in early diagnosis.

Through the employment of bioinformatics tools like Weighted Gene Co-Expression Network Analysis (WGCNA) in several studies, interactions between risk co-expressed genes have been identified in patients with SCZ (Torkamani et al. 2010; Kim et al. 2018) in addition to other mental diseases and healthy condition (Oldham et al. 2008). For instance, Wen et al. (Wen et al. 2020) identified 134 SCZ-specific key genes using WGCNA and Radulescu et al. (Radulescu et al. 2020) found 12 gene co-expression network modules in postmortem brain samples of SCZ patients. A number of other modules have been discovered using the construction of WGCNA in other studies (Liu et al. 2020; Zhang et al. 2020; Feltrin et al. 2019). To date, a number of susceptibility genes and characteristic genetic changes have been identified in association with increased risk of SCZ development. The affected genes have been found to mainly play role in the natural neurodevelopment which is believed to be affected in SCZ and other psychiatric disorders. These genes include immune function-related gene loci on chromosome 6p22.1 (Purcell et al. 2009), chromosome 22q11.2 deletion (Bassett et al. 2002), alleles of 5-hydroxytryptamine (5-HT) receptor gene, and dopamine D3 receptor gene (O'Donovan and Owen 1999). Studies have shown that thousands of single-nucleotide polymorphisms (SNPs) and copy number variants (CNVs) can explain parts of the genetic susceptibility to SCZ (Maric and Svrakic 2012). Currently known genetic mechanisms, however, have not been convincing in the independent elucidation of the SCZ pathophysiology (Pries et al. 2017). Same with many other diseases, epigenetic regulations along with genetic alterations including methylation, acetylation, phosphorylation, and SUMOylation on DNA bases or histone residues have been found to be involved in molecular mechanisms associated with SCZ.

To our knowledge based on the literature, protein networks and protein-miRNA networks have not been yet

identified in SCZ. In this study, we aimed to identify aberrantly expressed genes and miRNAs and associated cellular mechanisms among Iranian SCZ patients using gene co-expression network analyses. Retrieved hub miRNAs and genes were confirmed using quantitative reverse transcriptase polymerase chain reaction (qRT-PCR) and their diagnostic values were assessed through receiver operating characteristic (ROC) curve. The retrieved candidate miRNAs and genes suggested as potential biomarkers for SCZ patients.

## Materials and Methods

### Data Collection and Processing

Three expression arrays including Gene Expression Omnibus (GEO) series numbers GSE93987, GSE38485, and GSE54914 were obtained from the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>). According to the datasets, in GSE93987 individual pyramidal cells in dorsolateral prefrontal cortex layers 3 or 5 were captured by laser microdissection from 36 subjects diagnosed with SCZ or schizoaffective disorder and matched healthy controls. In GSE38485, the gene co-expression network in whole blood samples retrieved from 106 SCZ patients was compared to those from 96 healthy controls. In GSE54914 through a processing method, the probes were converted to the corresponding genes using their platforms, then the raw data were normalized by the quantile normalization function in the Limma package (Smyth 2005).

### Meta-analysis

Using the metaDE package (James et al. 2018), meta-analysis was performed for the two mentioned mRNA expression datasets, and differentially expressed genes (DEGs) between SCZ and healthy samples were identified. A false discovery rate (FDR) < 0.05 was selected as the significance threshold for screening of the DEGs.

### Weighted Gene Co-expression Network Construction and Module Detection

After identification of DEGs, the corresponding gene co-expression networks were constructed using the WGCNA package (Langfelder and Horvath 2008). For network construction, first a similarity matrix was made by calculating the correlations of all gene pairs. By applying the pickSoft-Threshold function, the appropriate soft-thresholding power  $B$  was selected to assess the scale-free topology. Afterward, to reduce the noise and spurious association impact, adjacency was transformed to the topological overlap matrix and

the corresponding dissimilarity was calculated. Hierarchical clustering was utilized to produce a dendrogram of genes. The modules with significant association with the measured clinical traits were selected according to two parameters including correlation and  $p$ -value in the next step. Then, we quantified the association of individual genes with our trait of interest (status) by defining Gene Significance (GS) as the absolute value of the correlation between the gene and the corresponding trait. For each module, we also defined a quantitative measure of module membership (MM) as the correlation between the module eigengene and the gene expression profile. This allows us to quantify the similarity of all array genes to each module. In the final step of network analysis, using the GS and MM measures we identified the genes with high significance for the status as well as high MM in the selected modules.

### Protein–Protein Interaction (PPI) Network Analysis and Pathway Enrichment Analysis

To identify the protein–protein interaction (PPI) information, the proteins of the selected modules were matched to the search tool for the retrieval of interacting genes (STRING) database (<https://string-db.org/cgi/input.pl>). The retrieved data was imported to the Cytoscape software (Shannon et al. 2003), and the corresponding PPI was visualized based on the degree and betweenness centrality. To elucidate the potential dysregulated signaling pathways, the ClueGO was utilized, and accordingly, the most related signaling pathways were identified (Bindea et al. 2009).

### Identification of Differentially Expressed miRNAs (DEMs)

By searching the GEO database, GSE54914 miRNA array was used for the identification of differentially expressed miRNAs (DEMs) between the SCZ patients and healthy controls. Significant miRNAs were identified based on  $p$ -value threshold of < 0.05 and  $\log_{2}FC > 1$ .

### MiRNA-Protein Interaction Network Construction and Identification of Hub miRNAs

DEMs and the most important genes retrieved from PPI were used for the construction of miRNA-protein interaction. For this purpose, the CyTargetLinker plugin (Kutmon et al. 2013) based on the Cytoscape software was employed, and a regulatory network between the retrieved miRNAs and genes was made using the miRBase (Griffiths-Jones et al. 2007), TargetScan (Agarwal et al. 2015), and TransmiR (Wang et al. 2010) databases. By consideration of the miRNA-protein network, three miRNAs and three genes were identified for further analyses.

## Validation of Detected miRNAs by the ROC Curve

The receiver operating characteristic (ROC) curve was plotted to confirm the diagnostic performance of the core miRNAs based on the GSE54914, and accordingly, the area under the curve (AUC) was estimated. The miRNAs with the  $p$ -value  $< 0.05$  and  $AUC > 0.8$  were considered as the strong potential diagnostic biomarkers.

## Validation of the Detected Genes and miRNAs Based on the Clinical Samples

To further confirmation of the results of bioinformatics data, qRT-PCR was conducted to evaluate the expression level of the selected genes and miRNAs in blood samples of the SCZ patients ( $n = 40$ ) recruited to Imam Khomeini, Mohammad Kermanshahi, and Farabi Hospitals, Kermanshah University of Medical Sciences, Kermanshah, Iran. Written informed consent was obtained from all patients. The demographic and clinical characteristics of the patients were provided in Supplementary Table 1.

## RNA Extraction and cDNA Production

Whole blood RNA purification kit (Qiagen Cat. No. 52304) was used for RNA extraction. Total RNA was extracted from 2-ml whole blood collected from each subject. Using reverse transcriptase enzyme, complementary DNA (cDNA) was synthesized, and then treated with DNase I enzyme to remove the genomic DNA. The purity of the extracted RNA was measured by calculation of the ratio  $A_{260}/A_{280}$  using UV spectrophotometry.

## Primer Design

The online miRNA designing tool (<http://genomics.dote.hu:8080/mirnadestool/>) was employed for the primer designing. To analyze the expression of the selected genes, primer design was performed using the GenScript online tool (<https://www.genscript.com/tools/real-time-pcr-tagman-primer-design-tool>)

## qRT-PCR

Primers arrived in the lyophilized form (Cinaclone, Iran). For preparation, sterile distilled water was added to each tube containing the lyophilized primer (based on the information provided for each primer), and the solution was placed in stoke at  $-20\text{ }^{\circ}\text{C}$ . Applied Biosystems Real-Time PCR instrument and Takara SYBR green kit were used for the analysis of miRNA and mRNA expression alterations. We also set up for each gene at each time, a negative control to examine the presence of contamination in each reaction.

The expression fold change of genes examined in this study was evaluated using the Threshold Cycle (CT) method by the formulas below. 5 s rRNA and GAPDH were used as the reference genes for this study.

$$R = 2^{-(\Delta\Delta\text{CT})}$$

$$\Delta\Delta\text{CT} = (\text{CT}_{\text{target}} - \text{CT}_{\text{reference}})_{\text{healthy}} - (\text{CT}_{\text{target}} - \text{CT}_{\text{reference}})_{\text{patient}}$$

## Results

### Meta-analysis

Two datasets with available SCZ mRNA expression data, including GSE93987 and GSE38485, and also 1 miRNA expression data, GSE54914, were downloaded from the GEO database. In the meta-analysis using the metaDE package, we identified 3840 DEGs, which were selected for further investigations (Fig. 1).


### Construction of Weighted Gene Co-expression Network

The WGCNA package was used in each mRNA expression datasets separately to construct the co-expressed networks and identify the co-expression modules retrieved from the results of the meta-analysis. Clustering dendrogram of the samples for the DEGs retrieved from each dataset was conducted based on their Euclidean distance (Figs. 2a and 3a). We then quantified the association between the individual genes and the trait of interest (SCZ and healthy) by defining the GS and MM values (Figs. 2b and 3b). A total number of 43 modules were identified for the DEGs retrieved from the GSE38485, and only genes of the lightcyan module were selected for further investigation. Seven modules were also identified for the GSE93987 and the blue module was selected for further analysis (Fig. 3b). According to the results, it is obvious that the GS and MM measures are highly correlated, illustrating that the genes with significant association with a trait are also the central elements of the modules associated with the trait (Figs. 2c and 3c). Based on Fig. 2c, genes of the light cyan module show correlation and  $p$ -value of 0.63 and  $3.5e^{-24}$ , respectively, and according to Fig. 3c, genes of the blue module demonstrate the corresponding values of 0.32 and  $4.6e^{-10}$ , respectively.

### Protein-Protein Interaction Network and Pathway Enrichment Analysis

All genes of the favorite modules (light cyan and blue) were used for the network construction. For the light cyan module,






**Fig. 1** Heatmap for DEGs between schizophrenia and healthy controls for two datasets including GSE93987 and GSE38485. Red colors represent upregulated genes and green colors represent downregulated genes

242 genes were selected, and using both STRING database and the Cytoscape software, the PPI network was made (Supplementary Fig. 1a). The color and size of each protein were set based on the degree and betweenness centrality. For the blue module, 362 proteins were selected, and according to the same protocol, the corresponding protein network was constructed (Supplementary Fig. 2a). To analyze the pathway enrichment, the CluGO plugin based on the Cytoscape software was used to illustrate the results of KEGG path

analysis for each of the gene groups separately (Supplementary Figs. 1b and 2b). For the genes of light cyan module, some cellular processes and signaling pathways were identified to be dysregulated. These included vitamin B6 binding, fucosyltransferase activity, single-stranded DNA binding, polyubiquitin modification-dependent protein binding, DNA replication origin binding, and double-stranded RNA binding. Also, dysregulated signaling pathways for the blue module included the epithelial cell signaling in



**Fig. 2** Network construction for GSE38485 using WGCNA. **(A)** Clustering dendrogram of samples based on their Euclidean distance. **(B)** Module-trait associations. Each row corresponds to a module eigengene, column to a trait. Each cell contains the correspond-

ing correlation and  $p$ -value. The table is color-coded by correlation according to the color legend. **(C)** A scatterplot of Gene Significance (GS) for weight vs. Module Membership (MM) in the light cyan module

*Helicobacter pylori* infection, vasopressin-regulated water reabsorption, dopaminergic synapse, proteasome, and *Salmonella* infection.


### Identification of the Differentially Expressed miRNAs

In the GSE54914 dataset, the miRNA expression profiles of 18 SCZ patients and 12 healthy controls were compared. Raw data were normalized via log<sub>2</sub> transformation and also quantile normalization function in the Limma package. The

results demonstrated 191 upregulated and 2 downregulated miRNAs (Fig. 4).

### MRNA-miRNA Network Construction

Based on two initial mRNA arrays, two miRNA-mRNA networks were made (Fig. 5A, B). According to the DEGs extracted from the GSE38485 and the DEMs identified by the GSE54914, a network was constructed (Fig. 6). As obvious, the hsa-miR-4429 shows direct effect on several DEGs including SORCS1, ZNF652, CHEK1, FAM63B,



**Fig. 3** Network construction for GSE93987 using WGCNA. **(A)** Clustering dendrogram of samples based on their Euclidean distance. **(B)** Module-trait associations. Each row corresponds to a module eigengene, column to a trait. Each cell contains the correspond-

ing correlation and  $p$ -value. The table is color-coded by correlation according to the color legend. **(C)** A scatterplot of Gene Significance (GS) for weight vs. Module Membership (MM) in the blue module

PDCD4, and FOXM1. Also, hsa-miR-574-5P directly regulates the expression of several other DEGs including MCM8, SLC35E1, HS6ST3, and ZRANB. The third miRNA hsa-miR-1827 was shown to regulate the expressions of some DEGs like ACBD7, XPNPEP3, PPM1K, and CENPM.

### Identification of Hub Genes

Using the DEGs retrieved from the GSE93987 and GSE38485 and DEMs identified by the GSE54914, two miRNA-mRNA networks were constructed (Fig. 5A, B). According to these networks, three miRNAs including hsa-miR-4429, hsa-miR-1827, and hsa-miR-574-5P play a central role. Therefore, we considered them as the hub miRNAs for further analyses. The common target genes of these miRNAs were identified using the Cytarargetlinker plugin based on the Cytoscape software (Fig. 6A, B). Several genes like CREBRF, ARPP19, TGFBR2, YWHAZ,

SRSF7, MAPK1IP1L, FAM117B, and BVES were demonstrated to be regulated by the hub miRNAs. We also evaluated the expression value of the detected genes in three independent datasets including the GSE62333, GSE93987, and GSE38485 (Fig. 6C).

### Validation of the Selected miRNAs by the ROC Curve

The ROC curves were plotted for the raw data retrieved from the results of Real-Time PCR and GSE54914 using a single-gene test in SCZ versus normal tissues for three miRNAs by plotting the sensitivity versus specificity. The results demonstrated significant values for the hub miRNAs. These included for hsa-miR-4429 (AUC: 0.88 and  $p$ -value: 0.0001, AUC: 0.9 and  $p$ -value: 0.0001), hsa-miR-1827 (AUC: 0.83 and  $p$ -value: 0.0001, AUC: 0.84 and  $p$ -value: 0.0013), and hsa-miR-574-5P (AUC: 0.76 and  $p$ -value: 0.0001, AUC: 0.9



and  $p$ -value: 0.0002). The AUC was analyzed by the Hanley and McNeil method (Supplementary Fig. 3).

### qRT-PCR

To evaluate the expression levels of the hub genes and miRNAs, qRT-PCR was carried out in 40 SCZ samples and matched healthy individuals (Fig. 7). Among the selected genes, CREBRF, TGFBR2, YWHAZ, and MAPK11P1L were shown to be downregulated in the SCZ samples compared to the controls, while the hub miRNAs were shown to be upregulated.


### Discussion

SCZ is one of the most debilitating mental disorders and a complex disease with an unknown etiology. Diagnosis is made based on the clinical manifestations on appearance, which makes the disease hardly manageable. Real-time diagnosis may help improve or slow down the progression of the consequences and response to treatment, which eventually promotes the patient's quality of life and social activities. Molecular diagnostics are potential nominates for on-time detection of susceptible or affected individuals. In accordance to the advancements in the past decades, numerous genetic studies have shown correlation between several genes and susceptibility to SCZ development in the people with familial history. Transcriptome studies also have demonstrated dysregulation of a growing number of miRNAs

in tissues and plasma of SCZ patients. MiRNAs as a novel group of ncRNAs act as the key regulators of gene expression playing substantial roles in several critical biological processes. Accordingly, dysregulation in their biogenesis has been associated with various human diseases including SCZ, for which a number of miRNAs are identified with aberrant expression in patient tissues compared with healthy individuals. Importantly, miRNAs are easily detected in human body fluids suggesting them as potential biomarkers for a wide variety of human disorders particularly in conditions for those no diagnostic approaches have been developed or diagnosed via invasive methods. Since accessibility to the brain biopsies is not possible for screening the SCZ susceptible individuals, development of a diagnostic approach based on the differentially expressed miRNAs potentially can benefit this society.


In the current study, we identified 3 hub miRNAs including hsa-miR-574-5P, hsa-miR-1827, and hsa-miR-4429 to be aberrantly expressed in blood sample of SCZ patients via network construction. The results were confirmed by the qRT-PCR. These miRNAs demonstrated acceptable diagnostic values in the ROC curve. The biological significance of these mRNAs and miRNAs has not been fully elucidated; however, they have been found to be dysregulated in several human disorders. Hsa-miR-574-5P, also known

**Fig. 4** Volcano plot display of differentially expressed miRNAs between schizophrenia and healthy controls. Every point in the plot represents a miRNA. Red points indicate miRNAs that are upregulated, and blue points indicate miRNAs that are downregulated










**Fig. 6** MiRNA-target interactions of the hub miRNAs including Has-miR-4429, Has-miR-1827, and HasmiR-574-5P. **(A)** Nodes between three big red nodes (Has-miR-4429, Has-miR-1827, and Has-miR-574-5P) are common target genes of them and were shown as yellow nodes. **(B)** Eight genes including CREBRF, ARPP19, TGFBR2,

YWHAZ, SRSF7, MAPK1IP1L, FAM117B, and BVES are common target genes of three mentioned hub genes. **(C)** The expression status of these common target genes evaluated in three independent expression arrays (GSE62333, GSE93987, and GSE38485) and the results are shown as heatmap

as miR-574-5P, has been studied as a malignancy promoter and cancer biomarker for several human tumors like lung cancer (Foss et al. 2011), nasopharyngeal carcinoma (Lin et al. 2020), colorectal cancer (Cui et al. 2014), and squamous cell carcinoma (Yang et al. 2013) in addition to its acceptable prognostic ability in prediction of prognosis in the patients with incident asthma (Li et al. 2021a). In lung cancer, vesicle-derived miR-574-5P is known to regulate prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) expression via Toll-like receptors (TLR) 7/8 (Donzelli et al. 2021). MiR-574-5P/TLR 7/8 axis is also identified to be involved in the pathogenesis of rheumatoid arthritis via induction of osteoclast differentiation (Hegewald et al. 2020). Additionally, it is known to be affected in several other human disorders such as rheumatoid arthritis (Hegewald et al. 2020), kidney injury in sepsis (Liu et al. 2021), and diabetes mellitus (Wang et al.

2021a). Regulatory functions of miR-574-5p also have been found on complement C7 involved in the pathogenesis of diabetic nephropathy (Guo et al. 2021), and serum lipids and blood glucose (Wang et al. 2021a) in addition to its antiviral activity against hepatitis B virus (HBV) (Wu et al. 2021). Another hub miRNA hsa-miR-1827 in addition to playing role in various human malignancies (Zhou et al. 2021a; Shen et al. 2021; Guo et al. 2020; Wang et al. 2020), regulation of osteogenic differentiation (Zhu et al. 2017), and primary immune thrombocytopenia (ITP) (Sun et al. 2021) has been found as a potential biomarker with dysregulation in samples of patients with Alzheimer's disease (Soleimani Zakeri et al. 2020). According to the literature, the third miRNA hsa-miR-4429 have been associated with particularly various types of cancer with tumor suppressor functions (Zhou et al. 2021b; Li et al. 2021b; Wang et al. 2021b), although in a few



**Fig. 7** Real-Time PCR analysis for miRNAs and mRNAs for schizophrenia in comparison with healthy controls in blood samples. Bar diagram shows fold change in expression of core miRNAs. Statistical

comparisons were made with respective 5 s rRNA for miRNAs and GAPDH for mRNAs.  $p < 0.05$  for all genes and miRNAs

studies is found to be dysregulated in other conditions like acute ischemic stroke and biliary atresia (Jickling et al. 2014; Dong et al. 2016). Among the DEGs, some have been known to play a role in neural health and function with changes contributing to neurological/neurodegenerative diseases like axon degeneration, Alzheimer's disease, and bipolar disease (Soleimani Zakeri et al. 2020; Li et al. 2021c; Reitz et al. 2011; Starnawska et al. 2016; Kim et al. 2001; Maycox et al. 2009). Taken together, the interaction we identified between the hub miRNAs and the hub genes can suggest potential role for the miRNAs in the SCZ pathogenesis. Still not elucidated, the association of these genes with the SCZ pathogenesis can be further explored to find diagnostic and therapeutical targets.

Overall, identification of these miRNAs and genes, already have been associated with several human diseases, in blood samples of SCZ patients may help develop a diagnostic approach via easily detection in the peripheral blood and so, earlier screening and monitoring the SCZ-affected individuals can be achieved. Further investigations are recommended to explore the biological functions and potential clinical application of these miRNAs.

In conclusion, this study is one of the first attempts to construct a miRNA-mRNA network using the metaDE, WGCNA, and limma packages. We identified 4 hub genes

and 3 miRNAs in three separate datasets. Then, their expression levels were verified by other independent datasets and real-time PCR, and their prognostic and diagnostic power was validated by the ROC curve.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s12031-021-01945-0>.

**Author Contribution** Conceptualization: Yazdan Rahmati and Omran Davarinejad; computational analyses and experiments: Yazdan Rahmati, Sajad Najafi, and Hossein Zhaleh; writing manuscript: Yazdan Rahmati, Sajad Najafi, Farzaneh Golmohammadia, Farnaz Radmehra, Mostafa Alikhania, and Reza Heidari Moghadam; and supervision and finalization: Yazdan Rahmati.

**Funding** This article is the result of the findings of the research project 97542 that was approved and financed by Vice Chancellor for Research and Research and Technology, Kermanshah University of Medical Sciences, Kermanshah, Iran.

**Availability of Data and Materials** The data used to support the findings of this study are available from the corresponding author upon request.

## Declarations

**Ethics Approval and Consent to Participate** All patients gave their signed written informed consent letters. Medical Research and Ethical Committee of Kermanshah University of Medical Sciences (Kermanshah,

Iran; registration no. IR.KUMS.REC.1397.490; grant number 97542) approved the study performed under ethical principles contained in the 7th and current (2013) editions of Helsinki Declaration.

**Conflict of Interest** The authors declare no competing interests.

## References

- Agarwal V, Bell GW, Nam JW, Bartel DP (2015) Predicting effective microRNA target sites in mammalian mRNAs. *Elife* 4:e05005
- Bassett AS, Chow EW, Weksberg R, Brzustowicz L (2002) Schizophrenia and genetics: new insights. *Curr Psychiatry Rep* 4(4):307–314
- Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A et al (2009) ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* 25(8):1091–1093
- Bushati N, Cohen SM (2007) microRNA functions. *Annu Rev Cell Dev Biol* 23(1):175–205
- Cui Z, Tang J, Chen J, Wang Z (2014) Hsa-miR-574-5p negatively regulates MACC-1 expression to suppress colorectal cancer liver metastasis. *Cancer Cell Int* 14(1):1–9
- Dong R, Shen Z, Zheng C, Chen G, Zheng S (2016) Serum microRNA microarray analysis identifies miR-4429 and miR-4689 are potential diagnostic biomarkers for biliary atresia. *Sci Rep* 6:21084
- Donzelli J, Proestler E, Riedel A, Nevermann S, Hertel B, Guenther A et al (2021) Small extracellular vesicle-derived miR-574-5p regulates PGE2-biosynthesis via TLR7/8 in lung cancer. *J Extracell Vesicles* 10(12):e12143-e
- Du Y, Tan WL, Chen L, Yang ZM, Li XS, Xue X et al (2021) Exosome transplantation from patients with schizophrenia causes schizophrenia-relevant behaviors in mice: an integrative multi-omics data analysis. *Schizophr Bull* 47(5):1288–1299
- Du Y, Yu Y, Hu Y, Li XW, Wei ZX, Pan RY et al (2019) Genome-wide, integrative analysis implicates exosome-derived MicroRNA dysregulation in schizophrenia. *Schizophr Bull* 45(6):1257–1266
- Feltrin ASA, Tahira AC, Simões SN, Brentani H, Martins DC Jr (2019) Assessment of complementarity of WGCNA and NERI results for identification of modules associated to schizophrenia spectrum disorders. *PLoS One* 14(1):e0210431-e
- Foss KM, Sima C, Ugolini D, Neri M, Allen KE, Weiss GJ (2011) miR-1254 and miR-574-5p: serum-based microRNA biomarkers for early-stage non-small cell lung cancer. *J Thorac Oncol* 6(3):482–488
- Griffiths-Jones S, Saini HK, Van Dongen S, Enright AJ (2007) miRBase: tools for microRNA genomics. *Nucleic Acids Res* 36(suppl\_1):D154–D158
- Guo X, Wang Z, Sun Q, Sun C, Hua H, Huang Q (2020) The inhibitory effect of microRNA-1827 on anoikis resistance in lung adenocarcinoma A549 cells via targeting caveolin-1. *Acta Biochim Biophys Sin* 52(10):1148–1155
- Guo H, Yan Z, Hu Y, Huang X, Pan C (2021) Complement C7 is specifically expressed in mesangial cells and is a potential diagnostic biomarker for diabetic nephropathy and is regulated by miR-494-3p and miR-574-5p. *Diabetes Metab Syndr Obes* 14:3077–3088
- Hauberg ME, Roussos P, Grove J, Børghlum AD, Mattheisen M, Schizophrenia Working Group of the Psychiatric Genomics Consortium (2016) Analyzing the role of MicroRNAs in schizophrenia in the context of common genetic risk variants. *JAMA Psychiat* 73(4):369–377
- He K, Guo C, He L, Shi Y (2017) MiRNAs of peripheral blood as the biomarker of schizophrenia. *Hereditas* 155(1):9
- Hegewald AB, Breitwieser K, Ottinger SM, Mobarrez F, Korotkova M, Rethi B et al (2020) Extracellular miR-574-5p induces osteoclast differentiation via TLR 7/8 in rheumatoid arthritis. *Front Immunol* 11:585282
- Islam MS, Khan MA, Murad MW, Karim M, Islam AB (2019) In silico analysis revealed Zika virus miRNAs associated with viral pathogenesis through alteration of host genes involved in immune response and neurological functions. *J Medic Virol* 91(9):1584–1594
- James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N et al (2018) Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* 392(10159):1789–1858
- Jickling GC, Ander BP, Zhan X, Noblett D, Stamova B, Liu D (2014) microRNA expression in peripheral blood cells following acute ischemic stroke and their predicted gene targets. *PLoS One* 9(6):e99283-e
- Kim Y, Giusti-Rodriguez P, Crowley JJ, Bryois J, Nonneman RJ, Ryan AK et al (2018) Comparative genomic evidence for the involvement of schizophrenia risk genes in antipsychotic effects. *Mol Psychiatry* 23(3):708–712
- Kim SH, Nairn AC, Cairns N, Lubec G (2001) Decreased levels of ARPP-19 and PKA in brains of Down syndrome and Alzheimer's disease. *J Neural Transm Suppl* 61:263–272
- Kloosterman WP, Plasterk RH (2006) The diverse functions of microRNAs in animal development and disease. *Dev Cell* 11(4):441–450
- Kuehner JN, Bruggeman EC, Wen Z, Yao B (2019) Epigenetic regulations in neuropsychiatric disorders. *Front Gen* 10(268)
- Kutmon M, Kelder T, Mandaviya P, Evelo CT, Coort SL (2013) CyTargetLinker: a cytoscape app to integrate regulatory interactions in network analysis. *PLoS One* 8(12):e82160
- Langfelder P, Horvath S (2008) WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* 9(1):1–13
- Leung AKL, Sharp PA (2010) MicroRNA functions in stress responses. *Mol Cell* 40(2):205–215
- Lichtenstein P, Yip BH, Björk C, Pawitan Y, Cannon TD, Sullivan PF et al (2009) Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *Lancet (london, England)* 373(9659):234–239
- Liu L, Zhao J, Chen Y, Feng R (2020) Metabolomics strategy assisted by transcriptomics analysis to identify biomarkers associated with schizophrenia. *Anal Chim Acta* 1140:18–29
- Liu S, Zhao L, Zhang L, Qiao L, Gao S (2021) Downregulation of miR-574-5p inhibits HK-2 cell viability and predicts the onset of acute kidney injury in sepsis patients. *Ren Fail* 43(1):942–948
- Lin Z, Chen M, Wan Y, Lei L, Ruan H (2020) miR-574-5p targets FOXN3 to regulate the invasion of nasopharyngeal carcinoma cells via Wnt/ $\beta$ -catenin pathway. *Technol Cancer Res Treat* 19:1533033820971659
- Li J, Tiwari A, Mirzakhani H, Wang AL, Kho AT, McGeachie MJ et al (2021a) Circulating microRNA: incident asthma prediction and vitamin D effect modification. *J Pers Med* 11(4):307
- Li W, Song Z, Jia N, Zhang C, Gao W, Wang L (2021b) microRNA-4429-5p suppresses the malignant development of colon cancer by targeting matrix metalloproteinase 16. *In Vitro Cell Dev Biol Anim* 57(7):715–725
- Li F, Lo TY, Miles L, Wang Q, Noristani HN, Li D et al (2021c) The Atr-Chek1 pathway inhibits axon regeneration in response to piezo-dependent mechanosensation. *Nat Commun* 12(1):3845
- Maes OC, An J, Sarojini H, Wang E (2008) Murine microRNAs implicated in liver functions and aging process. *Mech Ageing Dev* 129(9):534–541



- Mäki P, Veijola J, Jones PB, Murray GK, Koponen H, Tienari P et al (2005) Predictors of schizophrenia—a review. *Br Med Bull* 73–74(1):1–15
- Maric NP, Svrakic DM (2012) Why schizophrenia genetics needs epigenetics: a review. *Psychiatr Danub* 24(1):2–18
- Maycox PR, Kelly F, Taylor A, Bates S, Reid J, Logendra R et al (2009) Analysis of gene expression in two large schizophrenia cohorts identifies multiple changes associated with nerve terminal function. *Mol Psychiatry* 14(12):1083–1094
- Nelson PT, Wang WX, Rajeev BW (2008) MicroRNAs (miRNAs) in neurodegenerative diseases. *Brain Pathol* 18(1):130–138
- O'Donovan MC, Owen MJ (1999) Candidate-gene association studies of schizophrenia. *Am J Hum Genet* 65(3):587–592
- Oldham MC, Konopka G, Iwamoto K, Langfelder P, Kato T, Horvath S et al (2008) Functional organization of the transcriptome in human brain. *Nat Neurosci* 11(11):1271–1282
- Perkins DO, Jeffries CD, Jarskog LF, Thomson JM, Woods K, Newman MA et al (2007) microRNA expression in the prefrontal cortex of individuals with schizophrenia and schizoaffective disorder. *Genome Biol* 8(2):R27
- Pries L-K, Gülöksüz S, Kenis G (2017) DNA methylation in schizophrenia. In: Delgado-Morales R (ed) *Neuroepigenomics in Aging and Disease*. Springer International Publishing, Cham, pp 211–236
- Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF et al (2009) Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460(7256):748–752
- Radulescu E, Jaffe AE, Straub RE, Chen Q, Shin JH, Hyde TM et al (2020) Identification and prioritization of gene sets associated with schizophrenia risk by co-expression network analysis in human brain. *Mol Psychiatry* 25(4):791–804
- Reitz C, Tokuhiro S, Clark LN, Conrad C, Vonsattel J-P, Hazrati L-N et al (2011) SORCS1 alters amyloid precursor protein processing and variants may increase Alzheimer's disease risk. *Ann Neurol* 69(1):47–64
- Richetto J, Meyer U (2021) Epigenetic modifications in schizophrenia and related disorders: molecular scars of environmental exposures and source of phenotypic variability. *Biol Psychiat* 89(3):215–226
- Seven M, Karatas OF, Duz MB, Ozen M (2014) The role of miRNAs in cancer: from pathogenesis to therapeutic implications. *Future Oncol* 10(6):1027–1048
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D et al (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13(11):2498–2504
- Sharma RP, Grayson DR, Gavin DP (2008) Histone deacetylase 1 expression is increased in the prefrontal cortex of schizophrenia subjects: analysis of the National Brain Databank microarray collection. *Schizophr Res* 98(1–3):111–117
- Shen A, Tong X, Li H, Chu L, Jin X, Ma H et al (2021) TPPP3 inhibits the proliferation, invasion and migration of endometrial carcinoma targeted with miR-1827. *Clin Exp Pharmacol Physiol* 48(6):890–901
- Shi W, Du J, Qi Y, Liang G, Wang T, Li S et al (2012) Aberrant expression of serum miRNAs in schizophrenia. *J Psychiatr Res* 46(2):198–204
- Smyth GK (2005) *Limma: linear models for microarray data*. Springer, Bioinformatics and computational biology solutions using R and Bioconductor, pp 397–420
- Soleimani Zakeri NS, Pashazadeh S, MotieGhader H (2020) Gene biomarker discovery at different stages of Alzheimer using gene co-expression network approach. *Sci Rep* 10(1):12210
- Starnawska A, Demontis D, McQuillin A, O'Brien NL, Staunstrup NH, Mors O et al (2016) Hypomethylation of FAM63B in bipolar disorder patients. *Clin Epigenetics* 8(1):52
- Sun Y, Wang MJ, Cao XT, Liu WY, Chen HY, Ding XQ et al (2021) Expression of microRNAs in peripheral blood of patients with primary immune thrombocytopenia and its correlation with the imbalance of Th1/Th2 cell. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 29(5):1570–1576
- Tang X, Tang G, Özcan S (2008) Role of microRNAs in diabetes. *Biochim Et Biophys Acta (BBA)-Gene Reg Mech* 1779(11):697–701
- Taylor MA, Gaztanaga P, Abrams R (1974) Manic-depressive illness and acute schizophrenia: a clinical, family history, and treatment-response study. *Am J Psychiatry* 131(6):678–682
- Treiber T, Treiber N, Meister G (2012) Regulation of microRNA biogenesis and function. *Thromb Haemost* 107(04):605–610
- Torkamani A, Dean B, Schork NJ, Thomas EA (2010) Coexpression network analysis of neural tissue reveals perturbations in developmental processes in schizophrenia. *Genome Res* 20(4):403–412
- Van Wijnen AJ, Van De Peppel J, Van Leeuwen JP, Lian JB, Stein GS, Westendorf JJ et al (2013) MicroRNA functions in osteogenesis and dysfunctions in osteoporosis. *Curr Osteoporos Rep* 11(2):72–82
- van Os J, Kapur S (2009a) Schizophrenia. *Lancet (london, England)* 374(9690):635–645
- van Os J, Kapur S (2009b) Schizophrenia. *The Lancet* 374(9690):635–645
- Wang F, Li Z, Zhao M, Ye W, Wu H, Liao Q et al (2021a) Circulating miRNAs miR-574-5p and miR-3135b are potential metabolic regulators for serum lipids and blood glucose in gestational diabetes mellitus. *Gynecol Endocrinol* 37(7):665–671
- Wang J, Xie S, Liu J, Li T, Wang W, Xie Z (2021b) MicroRNA-4429 suppresses proliferation of prostate cancer cells by targeting distal-less homeobox 1 and inactivating the Wnt/ $\beta$ -catenin pathway. *BMC Urol* 21(1):40
- Wang J, Lu M, Qiu C, Cui Q (2010) TransmiR: a transcription factor–microRNA regulation database. *Nucleic Acid Res* 38(suppl\_1):D119–D122
- Wen Y-D, Xia Z-W, Li D-J, Cheng Q, Zhao Q, Cao H (2020) Genetic profiles playing opposite roles of pathogenesis in schizophrenia and glioma. *Journal of Oncology* 2020:3656841
- Wang Y, Gao R, Li J, Tang S, Li S, Tong Q et al (2020) Circular RNA hsa\_circ\_0003141 promotes tumorigenesis of hepatocellular carcinoma via a miR-1827/UBAP2 axis. *Aging* 12(10):9793–9806
- Wu W, Wu D, Yan W, Wang Y, You J, Wan X et al (2021) Interferon-induced macrophage-derived exosomes mediate antiviral activity against hepatitis B virus through miR-574-5p. *J Infect Dis* 223(4):686–698
- Yang M, Liu R, Sheng J, Liao J, Wang Y, Pan E et al (2013) Differential expression profiles of microRNAs as potential biomarkers for the early diagnosis of esophageal squamous cell carcinoma. *Oncol Rep* 29(1):169–176
- Zhang Y, You X, Li S, Long Q, Zhu Y, Teng Z et al (2020) Peripheral blood leukocyte RNA-Seq identifies a set of genes related to abnormal psychomotor behavior characteristics in patients with schizophrenia. *Medic Sci Moni: Int Med J Experiment Clin Res* 26:e922426-e
- Zhou Z, Zheng X, Mei X, Li W, Qi S, Deng Y et al (2021a) Hsa\_circ\_0080229 upregulates the expression of murine double minute-2 (MDM2) and promotes glioma tumorigenesis and invasion via the miR-1827 sponging mechanism. *Ann Translat Med* 9(9):762
- Zhou S, Qian K, Yu S, Zhao Y, Shen Q, Li Y (2021b) MiR-4429 alleviates malignant behaviors of lung adenocarcinoma through Wnt/ $\beta$ -catenin pathway. *Cancer Biother Radiopharma*
- Zhu S, Peng W, Li X, Weng J, Zhang X, Guo J et al (2017) miR-1827 inhibits osteogenic differentiation by targeting IGF1 in MSMSCs. *Sci Rep* 7:46136